

Toward a dynamical model for prime numbers

Claudio Bonanno* Mirko S. Mega†

Abstract

We show one possible dynamical approach to the study of the distribution of prime numbers. Our approach is based on two complexity methods, the Computable Information Content and the Entropy Information Gain, looking for analogies between the prime numbers and intermittency.

1 Introduction

Nowadays one of the most famous open problem in mathematics is the study of the distribution of prime numbers. This difficult problem has been studied in the last two centuries by many mathematicians, using many different techniques. More recently also in the physical literature it is possible to find many papers devoted to prime numbers, with two different points of view: a numerical approach using methods developed for physical systems; a theoretical approach, where prime numbers are found in some properties of physical systems.

This paper can be considered as a tentative to mix together the mathematical and the physical approach. We analyze prime numbers using numerical methods developed to study complexity in time series, and look for interpretations of the results in the realm of dynamical systems, in particular intermittent dynamical systems.

We first explain in Section 2 the complexity methods we use. Then in Section 3 we start to build a dynamical approach to prime numbers, looking for a dynamical model, and testing it by the complexity methods. It is interesting that the two methods seem to show some differences, sign of how difficult is to give a universal notion of complexity.

Finally in Section 4 we turn back to mathematics, to show that the prime numbers seem to have different statistical properties with respect to our dynamical models. Hence both the methods, that showed these discrepancies in different ways, seem to be able, if used together, to make a deep analysis of time series.

*Dipartimento di Matematica, Università di Pisa, via Buonarroti 2/a, 56127 Pisa (Italy), email: jbonanno@mail.dm.unipi.it;

†Dipartimento di Fisica, Università di Pisa, via Buonarroti 2, 56127 Pisa (Italy), email: jmstefano@math.uc3m.es;

2 Complexity methods for time series

There are many methods used in literature to study the “complexity” of a time series. In this paper we used two methods that follow from the interpretation of a time series as an information source: Computable Information Content ([1],[2]) and Diffusion Entropy ([3],[4]). Both methods aim to study the complexity of an underlying dynamics that produce the time series. The two methods have been studied also in [5].

2.1 Computable Information Content

Let (X, T, μ) be an invariant dynamical system, where X is a compact metric space (the phase space) with the Borel σ -algebra, $T : X \rightarrow X$ is a measurable transformation of X , and μ is a T -invariant measure on X , that is $\mu(T^{-1}(A)) = \mu(A)$ for all measurable sets $A \subseteq X$ (where $T^{-1}(A)$ is the counter-image of the set A).

One of the classical indicators of chaotic behavior of a dynamical system is given by the *Kolmogorov-Sinai (KS) entropy* $h_\mu(X, T)$, defined for probability T -invariant measures μ on X ([6],[7]). Chaitin and Kolmogorov independently introduced another method based on information theory to study chaotic systems, the *Algorithmic Information Content (AIC)* (see [8] and [9]). The two indicators of chaos are shown to coincide.

To study the information produced by a dynamical system, let Z be a finite partition of the phase space X into measurable sets $\{I_1, \dots, I_N\}$. To any point $x \in X$ we can associate an infinite string $\omega = (\omega_0, \omega_1, \dots, \omega_n, \dots)$, where for all $n \in \mathbb{N}$ it holds $\omega_n \in \mathcal{A} = \{1, \dots, N\}$ and $T^n(x) \in I_{\omega_n}$. The set $\mathcal{A}$ is called the *alphabet* associated to the partition Z , and the set of such strings ω is denoted by $\Omega \subseteq \mathcal{A}^{\mathbb{N}}$. We call the map $\varphi_Z : X \rightarrow \Omega$ the *symbolic representation* of (X, T) relative to Z . The space Ω is a compact metric space, with metric given by

$$d(\omega, \omega') = \sum_{n \in \mathbb{N}} \frac{1 - \delta(\omega_n, \omega'_n)}{2^n}$$

where $\delta(i, j) = 0$ if $i \neq j$ and $\delta(i, j) = 1$ if $i = j$. The Borel σ -algebra on Ω is generated by the so-called *cylinders* $C_\omega^{(k, h)}$, defined for all integers $k \leq h$ and $\omega \in \Omega$ by

$$C_\omega^{(k, h)} = \{\omega' \in \Omega / \omega'_n = \omega_n \ \forall \ n = k, \dots, h\}$$

Finally let $\sigma : \Omega \rightarrow \Omega$ be the continuous transformation of Ω given by $(\sigma(\omega))_n = \omega_{n+1}$ for all $n \in \mathbb{N}$. The map σ is called the *shift map*. Any T -invariant measure μ on X induces by the symbolic representation a σ -invariant measure $\nu := (\varphi_Z)_* \mu$ on Ω . Then (Ω, σ, ν) is a dynamical system, called the *symbolic dynamical system*.

When the symbolic representation φ_Z is injective then the dynamical systems (X, T, μ) and (Ω, σ, ν) are topologically conjugate. In this case the entropies $h_\mu(X, T, Z)$ and $h_\nu(\Omega, \sigma)$ coincide, for any probability measures μ and $\nu = (\varphi_Z)_* \mu$, respectively T and σ -invariant. Here $h_\mu(X, T, Z)$ is the KS entropy

of the dynamical system (X, T, μ) with respect to the partition Z . It holds $h_\mu(X, T) = \sup \{h_\mu(X, T, Z) \mid Z \text{ is a finite partition of } X\}$ and the supremum is attained on *generating* partitions.

The method of symbolic dynamics tells us that to study the information produced by the dynamical system (X, T, μ) , we can look at the different symbolic representations of the system, and in particular to those corresponding to generating partitions. Then we need to study the information in some sense contained in strings. One way is given by the *Algorithmic Information Content* (*AIC*).

Let s be a finite string on an alphabet $\mathcal{A}$, and let C be a universal Turing machine, then if p denotes a program given to C as input, we define

$$AIC(s) = \min \{|p| \mid C(p) = s\} \quad (1)$$

where $|p|$ is the binary length of the program p . In words, the *AIC* of a finite string s is the binary length of the smallest program p that gives s as output when run on a universal Turing machine.

The idea underlying the definition of the KS entropy is to estimate the rate with which the information necessary to reconstruct an orbit of the dynamical system increases with time. So, analogously, we study the rate with which the information contained in an infinite string increases with the length of the string. Formally, let $\omega \in \Omega$ be an infinite string, then the *complexity* $K(\omega)$ is defined as

$$K(\omega) = \limsup_{n \rightarrow \infty} \frac{AIC(\omega^n)}{n} \quad (2)$$

where $\omega^n = (\omega_0, \dots, \omega_{n-1})$ is the n -long string made by the first n symbols of the infinite string ω .

At this point it is immediate, by the method of symbolic representation of a dynamical system, to introduce a notion of complexity for orbits of a dynamical system using the AIC. Let indeed (X, T, μ) be an invariant dynamical system, and let Z be a finite measurable partition of X . Then via the map φ_Z to each point $x \in X$ we associate an infinite string $\omega \in \Omega \subseteq \mathcal{A}^\mathbb{N}$, where $\mathcal{A}$ is the finite alphabet associated to Z . Then the *complexity* $K(x, Z)$ of a point x with respect to the partition Z is given by

$$K(x, Z) := K(\varphi_Z(x)) = \limsup_{n \rightarrow \infty} \frac{AIC(x, n, Z)}{n} \quad (3)$$

where $AIC(x, n, Z) := AIC(\varphi_Z(x)^n)$.

The complexity of a point x and the KS entropy of a dynamical system are related by the following result (see [10],[2]): let (X, T, μ) be an invariant dynamical system, and let μ be a probability measure, then

$$\int_X K(x, Z) \, d\mu = h_\mu(X, T, Z) \quad (4)$$

for any finite measurable partition Z of X . In particular if the probability measure μ is also ergodic then

$$K(x, Z) = h_\mu(X, T, Z) \quad (5)$$

for μ -almost any $x \in X$. To obtain the KS entropy of a system, one can choose a generating partition Z and $h_\mu(X, T, Z) = h_\mu(X, T)$. If the invariant ergodic measure μ is infinite then (see [11] for exact hypothesis on systems with infinite measures)

$$K(x, Z) = 0 \quad (6)$$

for μ -almost any $x \in X$.

This result on complexity is not particular relevant for the case of probability invariant measures when $h_\mu(X, T) = 0$, but still the system is chaotic (these systems are called *weakly chaotic*), or for infinite invariant measures. In these cases what one can do is to look at the rate of increase of the *AIC* for the orbits of the system. Indeed, if $h_\mu(X, T, Z) > 0$ then, for a finite ergodic system, it holds $AIC(x, n, Z) \sim h_\mu(X, T, Z)n$ for μ -almost any x , but for ergodic weakly chaotic systems and for ergodic infinite systems it holds $AIC(x, n, Z) = o(n)$. Hence what one can do to classify the complexity of the system is to study the asymptotic behavior of the *AIC*. This has been done for some examples of weakly chaotic and infinite systems, in [2] the results are collected.

The problem with this approach is that the Algorithmic Information Content is a function that is not computable, in the sense that does not exist an algorithm able to compute the AIC for any finite string. This result is related to the Turing machine halting problem. Hence what one can do in practice is to study an algorithm which approximates the AIC. We used *compression algorithms*, that are formally defined in [2]. Given a finite string $s \in \mathcal{A}^*$, the set of all finite words in the alphabet $\mathcal{A}$, a compression algorithm takes it as input and gives as output a finite string $s' \in \{0, 1\}^*$, that is in some sense a string which contains all the information necessary to reconstruct s . We use as approximation of $AIC(s)$ the length of the compressed string s' . We call it *Computable Information Content (CIC)* of s . In the following we use a particular compression algorithm called *CASToRe* (introduced in [2]), that has been tested on some weakly chaotic and infinite systems, giving good approximations of the AIC. So we have $CIC(s) = |CASToRe(s)|$, where $|\cdot|$ denotes the length of a string.

2.2 Diffusion Entropy (DE)

The main idea of this method is to investigate the time series complexity building up a diffusion process, without any kind of pre-processing of the series. This technique is based principally on the Continuous time random walk (CTRW) and on the Generalized Central Limit Theorem (GCLT). It is very sensitive to time randomness, when the deviation from Poisson statistics generates Lévy diffusion, and to the trend of a non-stationary series, as in prime numbers. Let ξ_i , with $i = 1, \dots, M$, be a sequence of M numbers. The purpose of the DE is to establish the possible existence of a scaling, either normal or anomalous, in the most efficient way as possible, without altering the data with any form of detrending. Let l be a integer number, fitting the condition $1 \leq l \leq M$. This integer number will be referred as time. For any given l , there are $M - l + 1$

sub-sequence defined by

$$\xi_i^{(s)} \equiv \xi_{i+s} \quad s = 0, \dots, M - l \quad (7)$$

For any of this sub-sequence, a diffusion trajectory is built up, labeled with the index s , defined by

$$x^{(s)}(l) = \sum_{i=1}^l \xi_i^{(s)} \quad (8)$$

This position can be imagined as referring to a Brownian particle that, at regular intervals of time, has been jumping forward or backward, according to the prescription of the corresponding sub-sequence. This means that the particle, before reaching the position that it holds at time l , has been making l jumps. The jump made at the i -th step has the intensity $|\xi_i^{(s)}|$ and is forward or backward according to whether the number $\xi_i^{(s)}$ is positive or negative. Now the entropy of this diffusion process can be evaluated. To do that, the x -axis is partitioned into cells of size $\epsilon(l)$. When this partition is made, the cells are labeled. The number of how many particles are found in the same cell at a given time l is denoted by $N_i(l)$. This number is used to determine the probability that a particle can be found in the i -th cell at time l , $p_i(l)$, by means of

$$p_i(l) \equiv \frac{N_i(l)}{(M - l + 1)} \quad (9)$$

At this stage, the entropy of the diffusion process at time l is determined, and reads

$$S_{\epsilon(l)}(l) = - \sum_i p_i(l) \log[p_i(l)] \quad (10)$$

The easiest way to proceed with the choice of the cell size, $\epsilon(l)$, is to assume it independent of l and determine it by a suitable fraction of the square root of the variance of the fluctuation in the data ξ_i . A little comment on the way used to define the trajectories: the method is based on the idea of a moving window of size l that makes the s -th trajectory closely correlated to the next. The two trajectories have $l - 1$ values in common. A motivation of this choice is a possible connection with the Kolmogorov-Sinai (KS) entropy. The KS entropy of a symbolic sequence is evaluated by moving a window of size l along the sequence. Any window position corresponds to a given combination of symbols, and from the frequency of each combination, it is possible to derive the Shannon entropy $S(l)$. The KS entropy is given by the limit $\lim_{l \rightarrow \infty} S(l)/l$. Hence the idea is that the same sequence analyzed with the DE method, should yield at large values of l a well-defined scaling δ . As a simplifying assumption let consider large values of times to make the continuous assumption valid. In this case, the trajectories built up with the above illustrated procedure correspond to the following equation of motion:

$$\frac{dx}{dt} = \xi(t) \quad (11)$$

where $\xi(t)$ denotes the value of the time series under study at time t . This means that the function $\xi(l)$ is depicted as a function of t , thought of as a continuous time $t = l$. In this case, the Shannon entropy reads

$$S(x, t) = - \int_{-\infty}^{\infty} p(x, t) \log[p(x, t)] dx \quad (12)$$

being $p(x, t)$ the probability distribution function of the x variable at time t .

It has to be pointed out that scaling is a property implying that the probability distribution function can be read as

$$p(x, t) = \frac{1}{t^\delta} F\left(\frac{x}{t^\delta}\right) \quad (13)$$

Plugging equation (12) into equation (13), and after a little bit of algebra, $S(t)$ reads

$$S(x, t) = A + \delta \log(t) \quad (14)$$

where

$$A = \int_{-\infty}^{\infty} F(y) \log[F(y)] dy \quad (15)$$

It is now evident that this kind of techniques to detect scaling does not imply any form of detrending. This technique only works correctly in the case of a stationary time series, on the contrary, as can be analytically and numerically proved, if there is a trend we find a scaling with $\delta = 1$, as in the case of the prime numbers series. Whereas if the data ξ_i are the output of a process of independent and identically distributed random variables, then by the Central Limit Theorem the probability distribution function $p(x, t)$ converges to the Gauss distribution as $t \rightarrow \infty$, hence we find a scaling parameter $\delta = 0.5$.

On the same idea of not pre-processing the real data, a technique of conditional entropy, called *Entropy Information Gain (EIG)* [12] can be used, joined with the DE method, to detect the residual entropy associated to the independent fluctuations overlapped to the original trend. Let ξ_i be the original data and χ_i the time series produced using the hypothetical trend, then the corresponding trajectories are :

$$x^{(s)}(l) = \sum_{i=1}^l \xi_i^{(s)} \quad \text{and} \quad y^{(s)}(l) = \sum_{i=1}^l \chi_i^{(s)} \quad (16)$$

where we followed the same notations of equations (7) and (8). At this point we can repeat the same argument that we did for the DE to use continuous time, and define the EIG of the original data with respect to the hypothetical trend as the function $I(x, y, t)$ given by

$$I(x, y, t) = - \int_{-\infty}^{+\infty} p(y, t) dy \int_{-\infty}^{+\infty} p(x|y, t) \log p(x|y, t) dx \quad (17)$$

where $p(x, t)$ and $p(y, t)$ are the probability distribution functions associated to the trajectories x and y at time t , and $p(x|y, t) = p((x, y), t)/p(y, t)$, being

$p((x, y), t)$ the joint probability distribution function. After some simple algebra we find (see equation (12))

$$I(x, y, t) = S((x, y), t) - S(y, t) \quad (18)$$

that is an additive relation. Then it is evident that the better the hypothetical trend approximates the real time series, the smaller is the EIG. Moreover, applying the method of DE to the EIG, looking for scaling properties for the function $I(x, y, t)$, we interpret a $\delta = 0.5$ as a result of independent fluctuations in the original data, overlapped to the hypothetical trend.

3 Intermittency and the prime numbers

We now apply our methods to two different time series, studying the differences and the analogies. We first study the time series coming from an *intermittent* dynamical system, the *Manneville map*. The Manneville map is a dynamical system defined on the interval $I = [0, 1]$ by

$$T_z(x) = x + x^z \pmod{1} \quad z > 1 \quad (19)$$

where z is a real parameter. This map has the feature to have a fixed point in $x = 0$ which is not hyperbolic, indeed $T'_z(0) = 1$ for all $z > 1$. This makes the system interesting study, in particular for values of the parameter $z \geq 2$. Indeed in that range the invariant measure that is absolutely continuous with respect to the Lebesgue measure is infinite. For further properties we refer to [5],[13].

This map was originally introduced in [14] as a simple dynamical model for turbulence. Indeed the presence of the non-hyperbolic fixed point at the origin generates a phenomenon known as *intermittency*: an alternation of long laminar phases and short chaotic bursts. This intermittency is generated by the fact that when a point goes close to the origin it takes a long time to “escape” from it (the laminar phases), then it has a short period of time when it remains far from the origin (the chaotic burst), but then it falls again near the origin, and this process repeats on and on.

The behavior of the AIC for orbits of the Manneville map is studied in [13]. It is proved that,

$$\mathbb{E}[AIC(x, n, Z)] \sim \begin{cases} n & z < 2 \\ \frac{n}{\log n} & z = 2 \\ n^{\frac{1}{z-1}} & z > 2 \end{cases} \quad (20)$$

where $\mathbb{E}[\cdot]$ denotes the mean with respect to the Lebesgue measure and it has been chosen the partition $Z = \{[0, \tilde{x}), [\tilde{x}, 1]\}$, being $\tilde{x}$ the real number in $(0, 1)$ such that $x + x^z = 1$. The same behavior is obtained for the Computable Information Content $\mathbb{E}[CIC(x, n, Z)]$ using *CASToRe* ([2]).

The second time series we study is obtained from the series of prime numbers. Let $\{1, 2, 3, \dots\}$ be the integer numbers, we associate to each number the symbol

“1” if it is a prime number, and the symbol “0” if it is not. So we construct an infinite string $\bar{\omega} \in \{0, 1\}^{\mathbb{N}}$, whose first symbols are $\bar{\omega} = (0, 1, 1, 0, 1, 0, 1, \dots)$. The most famous question on prime numbers is “how many prime numbers are there in the first n integers?”. This question becomes in our formulation “how many “1”s are there in the first n symbols of $\bar{\omega}$?”. We refer to [15] for the formal approach. The first answer to this question was given by Gauss in 1849, and his result is contained in the

Prime Numbers Theorem. If $\pi(n)$ is the number of primes up to the first n integers, it holds

$$\pi(n) \sim \text{Li}(n) := \int_2^n \frac{dt}{\log t} \sim \frac{n}{\log n}$$

We now study what happens if we want to study the information content of the string $\bar{\omega}$. Since there are many algorithms to generate the prime numbers (think for example to Eratosthenes’ sieve), if we want to estimate $AIC(\bar{\omega}^n)$ it is clear that it is of the order $\log n$, since the algorithm that generates the primes only needs to know at what integer n it has to stop (this is done with $\log n$ bits of information). So the shortest program has length of the order $\log n$. But we imagine to forget the origin of the string $\bar{\omega}$ and look for its information content.

In [13] it is shown that the compression of a binary string done simply remembering how many “0”s there are between two consecutive “1”s, is the “best” compression for strings of the Manneville map (“best” in the sense that it approximates the behavior of the AIC up to a constant). This fact is due to the property of strings from the Manneville map that there are long sequences made only of “0”s (correspondent to the laminar phases) and short sequences with chaotic alternation of “0”s and “1”s (the chaotic bursts), and to the property that the passage through a chaotic burst can be considered as a phenomenon deleting the memory of the system, so what happens after a symbol “1” does not depend on what happened before. The string $\bar{\omega}$ of prime numbers has the same property of long sequences of “0” and rare appearances of “1”s, so we could try to apply this kind of compression to $\bar{\omega}$ to see what happens, even if the property of loss of memory is false for prime numbers. It is not difficult to see that the length of the new string obtained is related to the number of symbols “1” in the original string. So it is related to the function $\pi(n)$. This argument shows that we could imagine the string $\bar{\omega}$ as given from one of the orbits of the Manneville map with $z = 2$, both having the same asymptotic behavior of the information content (see equation (20)).

We remark that this approach to study prime numbers is similar to the probabilistic approach introduced by Cramér in [16], that is we assume that the string $\bar{\omega}$ is one of a family of strings, on which there is a probability measure μ . If then a property holds for μ -almost all strings of the family, then there is a “good” chance that it holds also for $\bar{\omega}$. Of course it is not formally true, the string $\bar{\omega}$ could be in the set of null μ -measure for which the property does not hold, but certainly it is a good first guess for the properties of $\bar{\omega}$. Our aim is to look for a dynamical system (X, T) whose orbits have (in some probabilistic sense)

the same statistical properties of $\bar{\omega}$, so that we can think of $\bar{\omega}$ as the symbolic representation of an orbit of the system, and guess some new properties of $\bar{\omega}$ from those of the system.

Following the idea explained above, we compare the statistical properties of strings generated by the Manneville map T_2 and the infinite string $\bar{\omega}$. We apply the methods introduced in Section 2 to study the complexity of the time series.

We first apply the compression algorithm *CASToRe*. We built the first 10^7 symbols of the string $\bar{\omega}$ and 10 orbits of the Manneville map T_2 of the same length, corresponding to different initial conditions (this because we remark that for the AIC of T_2 we only have results in mean with respect to the Lebesgue measure). In Figure 1 it is shown the behavior of the information content for all the series.




Figure 1: *The information content measured using CASToRe for the series of the prime numbers and ten different orbits of the Manneville map T_2 . The scale is bi-logarithmic.*

The graph is in bi-logarithmic scale, hence the behavior $CIC(n) \sim \frac{n}{\log n}$ becomes a straight line with angular coefficient 1, perturbed by a logarithm, that is smaller than $7 \log 10$. In this graph it is difficult to distinguish the different curves, but there is one that is practically straight, the one of $\bar{\omega}$. Of course the algorithm *CASToRe* is only an approximation of the AIC of our series, hence we cannot expect the curves to be exactly of the order $\frac{n}{\log n}$, but they

are approximations to it. However our aim is not to estimate the information content of the series, but simply to make a comparison between their behaviors, hence we can neglect the effects due to the use of the particular compression algorithm since they are the same on each series.

From this first figure we could then conclude that the Manneville map T_2 is a “good” dynamical system to generate the series of the prime numbers, in the sense explained above. However if we look at the same graphs in bi-linear scale (Figure 2), we notice a small difference between the straight line (the information content of $\bar{\omega}$) and the other curves.




Figure 2: *The same as in Figure 1, but in bi-linear scale.*

This small difference is not visible in the bi-logarithmic figure, since the bi-logarithmic scale reduces the differences at high values of n . Hence the information content of the string $\bar{\omega}$ seems to be asymptotically slightly different from the mean information content of orbits of T_2 .

This difference can be explained in terms of the approximation done in the Prime Numbers Theorem. Indeed the function $\text{Li}(n)$ can be estimated using the method of integration by parts, obtaining the approximation

$$\text{Li}(n) \sim \frac{n}{\log n} + \frac{n}{\log^2 n} + \mathcal{O}\left(\frac{n}{\log^3 n}\right)$$

and this shows that the number of primes in the first n integers (that is the

number of symbols “1” in the first n symbols of the string $\bar{\omega}$) grows slightly faster than $\frac{n}{\log n}$.

Then we apply the Entropy Information Gain method to the same series. First of all we study how good is the approximation of the function $\pi(n)$ given in the Prime Numbers Theorem. We build up the different trajectories to analyze just by the sum of the number of primes contained in the strings of length l . We use for the hypothetical trend the first approximation of the Prime Numbers Theorem, $\pi(n) \sim \frac{n}{\log n}$. In Figure 3 it is shown the behavior of the residual Shannon entropy, that follows a straight line with angular coefficient ≈ 0.4 .




Figure 3: *EIG for the prime numbers series, using as trend the first approximation of the function $Li(n)$.*

This value of the δ , close to the Gaussian value 0.5, seems to suggest that all the information of the primes series is contained in the Prime Number Theorem. To better investigate about the primes trend, we have also calculated the EIG with more terms approximating the function $Li(n)$, Figure 4, but the results doesn't change. This equality could be a limit for the sensibility of this method, that seems to feel only the first approximation of a hypothetical trend.

At this point we have calculated the EIG using as data the prime numbers series and as hypothetical trend the series of orbits produced by the Manneville




Figure 4: *The same as in Fig 3 but with the first two orders of approximation for $Li(n)$.*

Map for $z = 2$, Figure 5. Also in this case, the angular coefficient of the asymptotic straight line is ≈ 0.5 , due to the fact that in first approximation, this Manneville map contains the same information of the Prime Numbers Theorem, even if the compression method has shown some differences, that if coming from the approximation of the function $Li(n)$ are stationary, so cannot be interpreted as independent fluctuations on the trend.

So far we have applied theoretical results from number theory and from the theory of information content in intermittent dynamical systems, obtaining a “close” relationship between the prime number series $\bar{\omega}$ and the orbits of the Manneville map T_2 . To find a better dynamical model for the prime numbers, in the sense explained above, we could slightly perturb the map T_2 and see what happens for the information content. This is what is done below based only on numerical experiments done using the compression algorithm *CASToRe*.

The statistical properties of the maps T_z are induced by the behavior of the maps near the origin (the non-hyperbolic fixed point). Hence we have to perturb T_2 changing its behavior near the origin. We have chosen the following family of maps: given two positive real numbers a, b , the maps $T_{(a,b)}$ are defined by

$$T_{(a,b)}(x) = x + x^2 + ax^b \quad x \in (0, \tilde{x}) \quad (21)$$

where $\tilde{x}$ satisfies $\tilde{x} + \tilde{x}^2 + a\tilde{x}^b = 1$, and $T_{(a,b)}(x)$ is randomly chosen in $(0, 1)$




Figure 5: *EIG between the prime numbers and the different Manneville maps. The $\delta = 0.53$ correspond to the T_2 map, while the $\delta = 0.33$ to the new hybrid maps.*

according to the Lebesgue measure when $x > \tilde{x}$.

In Figures 6-9 the results of the experiments are shown. Each figure is in bi-linear scale, and there are plotted the curves corresponding to the information content of the string $\bar{\omega}$ (the straight line) and of some different orbits of the maps $T_{(a,b)}$, for (a,b) equal respectively to $(1,3)$, $(0.1,2)$, $(0.5,2)$ and $(1,2)$. From the figures, we obtain as a “good” guess for a dynamical model of the series of prime numbers the map $T_{(0.5,2)}$, even if in this case the results show a big difference for the behavior of the information content of different orbits (remember we have only results in mean for intermittent maps with an infinite invariant measure).

On the same line, we have calculated the EIG between the prime numbers series and the different perturbed Manneville maps $T_{(a,b)}$. The results are shown in Figure 5: in this figure there is the EIG result for T_2 , with $\delta = 0.53$, whereas all the perturbed maps give a $\delta = 0.33$, sign of an anti-correlation behavior for the residual Shannon entropy. Hence it seems that now the perturbation to the Manneville map gives better results from the point of view of compression algorithms, but the results seem worse from the point of view of the EIG.




Figure 6: *The information content measured using CASToRe for the series of the prime numbers and ten different orbits of the map $T_{(1,3)}$. The scale is bi-linear.*

4 Waiting times and distance between primes

In the analysis of the statistical behavior of time series, an important feature is the distance between two different events. In the orbits of intermittent dynamical systems the events are the chaotic bursts. The importance of these events for the intermittent systems is reflected by the fact that the “best” compression was done by simply recording how much time an orbit spends in the laminar zone (how many symbols “0” there are in the series) between two different chaotic bursts (corresponding to the appearance of the symbol “1”). Analogously, for the series of prime numbers the events are the appearance of the symbol “1”, that is the fact that an integer is prime. Hence the distance between two different events is the distance between two consecutive prime numbers. The series of the distances between consecutive prime numbers has been analyzed with different complexity methods, see for example [17], [18] and references therein.

In this section we compare the behavior of the distances between events for the prime numbers and the intermittent dynamical systems $T_{(a,2)}$.

For prime numbers there are many different results about the growth of the distance $(p_{n+1} - p_n)$ between consecutive primes. Using the probabilistic




Figure 7: *The same as in Figure 6, but for the map $T_{(0.1,2)}$.*

approach, Cramér ([16]) gave the conjecture

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log^2 p_n} = 1 \quad (22)$$

To investigate this relation, we have built the distance time series from the prime numbers series and we have applied the EIG method. In Figure 10 we shown as once more, the hypothetical trend for the distances between primes, given by Cramér conjecture, seems to contain most part of the information, yielding a $\delta = 0.5$, sign of independent fluctuations.

This result can be rephrased in terms of the time t_r needed to have r consecutive “0”s in $\bar{\omega}$. Indeed it holds that $t_r(\bar{\omega}) \geq \exp(\sqrt{r})$.

The analysis of the same quantity t_r was done in [19] for the family of maps $T_{(a,2)}$. The result is that for r big enough

$$K^{-1} \leq \frac{\mathbb{E}_\rho[t_r]}{r \log r} \leq K \quad (23)$$

for some positive constant K , where the mean $\mathbb{E}_\rho$ is done with respect to a probability measure ρ , absolutely continuous with respect to the Lebesgue measure.




Figure 8: *The same as in Figure 6, but for the map $T_{(0.5,2)}$.*

Hence if from the point of view of the information content of the orbits, the intermittent map $T_{(0.5,2)}$ seems to have, in mean with respect to the Lebesgue measure, the same behavior as $\bar{\omega}$, the results on the distances between consecutive events show big differences between the two different kinds of intermittency, as was suggested by the EIG results. If we also notice that the family of Manneville maps T_z has a phase transition at $z = 2$, we can conclude that the prime numbers are well approximated by an intermittent dynamical system similar to Manneville maps only to some extent.

5 Conclusion

We have shown the behavior of two complexity methods, Computable Information Content and Entropy Information Gain, on the prime numbers series, with an open eye on possible relations with dynamical systems, using theoretical and numerical results.

This paper is supposed only as a tentative to open a different approach to study prime numbers, the dynamical approach. Showing at the same time, the importance of different complexity notions, that on time series can show analogies, but also deep differences, as in this case.

Moreover we remark that in our approach we looked for a deterministic




Figure 9: *The same as in Figure 6, but for the map $T_{(1,2)}$.*

model for prime numbers, where the model is supposed to be fixed. A different approach, similar to the spirit of the analysis in [18], is to consider a model not fixed, but with characteristics varying with time. Our idea is however that to guess how the characteristics of the model has to vary with time is a problem of the same difficulty of the approximation of the function $\pi(n)$ of the Prime Numbers Theorem.

References

- [1] F.Argenti, V.Benci, P.Cerrai, A.Cordelli, S.Galatolo, G.Menconi, *Information and dynamical systems: a concrete measurement on sporadic dynamics*, Chaos Solitons Fractals **13** (2002), no.3, 461–469
- [2] V.Benci, C.Bonanno, S.Galatolo, G.Menconi, M.Virgilio, *Dynamical systems and computable information*, Discrete and Continuous Dynamical Systems - B, to appear (2003)
- [3] N.Scafetta, P.Hamilton, P.Grigolini, *The thermodynamics of social processes: the teen birth phenomenon*, Fractals **9** (2001), 193–208




Figure 10: *EIG between the prime numbers distances and the hypothetical trend $\log^2(p_n)$.*

- [4] P.Grigolini, L.Palatella, G.Raffaelli, *Asymmetric anomalous diffusion: an efficient way to detect memory in time series*, Fractals **9** (2001), 439–449
- [5] P.Allegri, V.Benci, P.Grigolini, P.Hamilton, M.Ignaccolo, G.Menconi, L.Palatella, G.Raffaelli, N.Scafetta, M.Virgilio, J.Yang, *Compression and diffusion: a joint approach to detect complexity*, Chaos Solitons Fractals **15** (2003), 517–535
- [6] C.Beck, F.Schlögl, “Thermodynamics of Chaotic Systems”, Cambridge University Press, 1993
- [7] J.R.Dorfam, “An Introduction to Chaos in Non Equilibrium Statical Mechanics”, Cambridge University Press, 1999
- [8] G.J.Chaitin, “Information, Randomness and Incompleteness. Papers on Algorithmic Information Theory”, World Scientific, 1987
- [9] A.N.Kolmogorov, *Combinatorial foundations of information theory and the calculus of probabilities*, Russ. Math. Surv. **38** (1983), 29–40
- [10] A.A.Brudno, *Entropy and the complexity of the trajectories of a dynamical system*, Trans. Moscow Math. Soc. **2** (1983), 127–151

- [11] C.Bonanno, “Applications of Information Measures to Chaotic Dynamical Systems”, Ph.D. thesis, University of Pisa, 2003
- [12] C.Shannon, W.Weaver, “Théorie mathématique de la communication”, Bibliothèque du CELP, 1975
- [13] C.Bonanno, S.Galatolo, *Orbit complexity for the Manneville maps*, work in preparation
- [14] P.Manneville, *Intermittency, self-similarity and $1/f$ spectrum in dissipative dynamical systems*, J. Physique **41** (1980), 1235–1243
- [15] T.M.Apostol, “Introduction to Analytic Number Theory”, Springer Verlag, 1976
- [16] H.Cramér, *Prime numbers and probability*, Skand. Mat.-Kongr. **8** (1935), 107–115
- [17] P.Kumar, P.Ch.Ivanov, H.E.Stanley, *Information entropy and correlations in prime numbers*, cond-mat/0303110 (2003)
- [18] N.Scafetta, T.Imholt, J.A.Roberts, B.J.West, *An intensity-expansion method to treat non-stationary time series: an application to the distance between prime numbers*, Chaos Solitons Fractals, this issue
- [19] M.Campanino, S.Isola, *Statistical properties of long return times in type I intermittency*, Forum Math. **7** (1995), 331–348